

Regional Characterisation of a Major Storage System: Gippsland Basin, Southeast Australia

Catherine M. Gibson-Poole^{1,2}, L. Svendsen^{1,2}, J. Underschultz^{1,3}, M. Watson^{1,2}, J. Ennis-King^{1,3}, P. van Ruth^{1,2}, E. Nelson², R. Daniel^{1,2} and Y. Cinar^{1,4}

¹CRC for Greenhouse Gas Technologies

²Australian School of Petroleum, The University of Adelaide

³CSIRO Petroleum

⁴The University of New South Wales

CO₂SC 2006 – Berkeley, CA – 20th March 2006



UNSW



ASP

Australian School of Petroleum



THE UNIVERSITY
OF ADELAIDE
AUSTRALIA



Presentation Outline

- Introduction
- Workflow for detailed site characterisation
- Gippsland Basin, southeast Australia
 - Base regional seal migration pathways
 - Kingfish Field: sequence stratigraphy and depositional model
 - Injectivity: reservoir quality, geometry & connectivity
 - Containment: seal capacity, migration pathways, trap mechanism, geomechanical assessment, hydrodynamic analysis
 - Capacity: 3D geological model & pore volume
 - Numerical flow simulation
- Conclusions



UNSW




Australian School of Petroleum



THE UNIVERSITY
OF ADELAIDE
AUSTRALIA



Introduction




Amount of CO₂ geologically stored influenced by:

- Rate of CO₂ migration
- Style of multiphase flow
- Rate of CO₂ dissolution
- Rate of chemical reaction with minerals

Controlled by many variables, including:

- Reservoir and seal structure
- Stratigraphic architecture
- Reservoir heterogeneity
- Faults/fractures
- Pressure/temperature conditions
- Hydrodynamics and chemistry of *in situ* formation fluids

Site Characterisation Workflow



UNSW



CSIRO



Australian School of Petroleum

ASP




THE UNIVERSITY
OF ADELAIDE
AUSTRALIA



Gippsland Basin, Southeast Australia

The Problem:

- New brown coal developments in Latrobe Valley, Victoria
- CO₂ emissions up to 50 Million tonnes/year



Potential Solution:

- Offshore Gippsland Basin
- Existing oil and gas fields (once depleted)
- Deeper saline formations



UNSW



CSIRO




Australian School of Petroleum



THE UNIVERSITY
OF ADELAIDE
AUSTRALIA



Buoyancy migration pathways at base regional seal




Selected Site Scenarios

Injection Scenarios:

Injection at several sites along regional migration pathways, sequentially & simultaneously, ramping up volume to 50 Mt/y

1. Kingfish Field: 15 Mt/y for 40 years
2. Fortescue Field: 15 Mt/y for 40 years
3. Basin centre & northern gas fields: 20 Mt/y for 40 years



UNSW



CSIRO



Australian School of Petroleum




THE UNIVERSITY OF ADELAIDE AUSTRALIA




Detailed Characterisation: Kingfish Field

Location map of Gippsland Basin



(modified after Power et al., 2001)

Stratigraphic column



(after Bernecker & Partridge, 2001)



UNSW



CSIRO



Australian School of Petroleum




THE UNIVERSITY OF ADELAIDE AUSTRALIA



Sequence Stratigraphy

Seismic Cross-Section (line G92A-3074A) between Kingfish-3 and Roundhead-1



UNSW




Australian School of Petroleum



THE UNIVERSITY
OF ADELAIDE
AUSTRALIA



Injectivity: Reservoir Quality



Thin section micrograph: Kingfish Fm, nearshore facies


Kingfish Fm

- 15-30 % porosity
- 10-10,000 mD perm
- Good to excellent reservoir quality


Geochemical reactions

- Reservoir units lack minerals reactive to CO_2
- Injectivity unlikely to be compromised

Containment: Seal Capacity (MICP analysis)



Containment: Migration Pathways




Intra-Latrobe Gp


- Stratigraphy dips down to west
- CO₂ will migrate updip to east

Top Latrobe Gp

- Base regional seal dips down to east
- CO₂ will migrate updip to west (towards Bream)



Containment: Migration Pathways Concept




Importance of intraformational baffles:

- increase length of CO₂ migration pathway
 - increase volume of pore space moved through
- = greater residual gas trapping & dissolution

Containment: Geochemical Trapping

Ideal Reservoir System



Lakes Entrance Fm

- High seal capacity
- Quartz & illitic-smectite
- = limited mineral reactions

Gurnard Fm


- Low permeability
- Calcium, iron & magnesium-bearing minerals

= significant potential for mineral trapping of CO₂

Kingfish Fm

- Moderate to high permeability
- Non-reactive minerals
- = limited mineral reactions

Containment: Geomechanics




Seismically-resolvable faults

- 3 cut the top Latrobe unconf.
- 7 terminate within Latrobe Gp
- Most have moderate to high fault reactivation potential
- However, most not in immediate migration pathway

Containment: Hydrodynamics

Post-production hydraulic head – influence short-term CO₂ fate (10s – 100s yrs)

Data from the mine site area is taken from "Latrobe Valley regional groundwater and land surface monitoring report five year review as at June 2000". #1000/8505/99 Geo-Eng Pty Ltd.



Capacity

CO₂ Storage Capacity

- Available pore volume calculated geologically
- Numerical simulation required to verify pore volume used (sweep efficiency)
- Sweep efficiency dependent on: rate of CO₂ migration, dissolution into formation water, precipitation of new minerals, fill-to-spill structural closures along migration path

Kingfish Field

- Calculated structural closure capacity (existing oil zone) and deeper intra-Latrobe stratigraphy
- Combined capacity > 600 Mt (sufficient for 15 Mt/y for 40 years)
- Intra-Latrobe stratigraphy 3 times the capacity of the structural closure – demonstrates how a deeper injection strategy may provide significantly more CO₂ storage capacity



UNSW



Australian School of Petroleum


ASP



THE UNIVERSITY
OF ADELAIDE
AUSTRALIA



Numerical Flow Simulation



UNSW




Australian School of Petroleum

ASP




THE UNIVERSITY
OF ADELAIDE
AUSTRALIA






- 15 Mt/y for 40 years
- Post-injection small shales 0–40 yrs



- 15 Mt/y for 40 years
- Post-injection small shales 40–400 yrs



Simulation results:

- Injection rate achievable – lower permeability or extensive shale barriers require more wells
- Migration time to the oil-bearing zone is 40-200 years for deep injection – less for shallow injection, more for wider shale barriers
- Storage capacity sufficient with deep injection – more CO₂ trapped as residual gas

- 15 Mt/y for 40 years
- Post-injection small shales 400–1140 yrs

Conclusions

Suitability of Kingfish Field/Gippsland Basin as CO₂ geological storage site:

- Complex stratigraphic architecture which slows vertical migration and increases residual gas trapping
- Non-reactive reservoir units with high injectivity
- Geochemically-reactive, low permeability reservoir just below regional seal to provide additional mineral trapping
- Several depleted oil fields to provide storage capacity coupled with transient flow regime that enhances containment
- Long migration pathways beneath competent regional seal
- Kingfish Field, in conjunction with other sites (e.g. Fortescue, northern gas fields), indicate that Gippsland Basin has sufficient capacity to store very large volumes of CO₂.

Site characterisation for CO₂ geological storage:

- Geological variability requires sites to be independently assessed, although similar workflow can be applied (i.e. injectivity, containment, capacity)
- Best addressed by multidisciplinary approach (i.e. detailed geoscience, engineering, economics and risk assessment) to provide integrated and comprehensive site characterisation



UNSW



Australian School of Petroleum



THE UNIVERSITY
OF ADELAIDE
AUSTRALIA



Acknowledgements: CO2CRC Participants



Australian Government

Geoscience Australia

Australian Greenhouse Office

Department of Industry, Tourism and Resources



**RIO
TINTO**



**THE UNIVERSITY
OF ADELAIDE
AUSTRALIA**



**STANWELL
CORPORATION LIMITED**



Schlumberger



curtin

University of Technology
Western Australia



MONASH University

ACARP

Australian Coal Association Research Program



UNSW



**THE UNIVERSITY OF
MELBOURNE**



SOLID ENERGY
Coals of New Zealand



Supporting participants: **Australian National University** | **CANSYD** | **Meiji University** |
| **The Process Group** | **University of Queensland** | **Whistler Research** |



UNSW



Australian School of Petroleum

ASP



**THE UNIVERSITY
OF ADELAIDE
AUSTRALIA**

